

Physical properties of $z \sim 4$ LBGs: differences between galaxies with and without $\text{Ly}\alpha$ emission

L.Pentericci¹, A. Grazian¹, A. Fontana¹, S. Salimbeni¹, P. Santini¹, C. De Santis¹, S. Gallozzi¹, and E. Giallongo¹

INAF - Osservatorio Astronomico di Roma, Via Frascati 33, I-00040, Monteporzio, Italy

ABSTRACT

Aims. We have analysed the physical properties of $z \sim 4$ Lyman Break Galaxies observed in the GOODS-S survey, in order to investigate the possible differences between galaxies where the $\text{Ly}\alpha$ is present in emission, and those where the line is absent or in absorption.

Methods. The objects have been selected from their optical color and then spectroscopically confirmed by Vanzella et al. (2005). From the public spectra we assessed the nature of the $\text{Ly}\alpha$ emission and divided the sample into galaxies with $\text{Ly}\alpha$ in emission and objects without $\text{Ly}\alpha$ line (i.e. either absent or in absorption). We have then used the complete photometry, from U band to mid infrared from the GOODS-MUSIC database, to study the observational properties of the galaxies, such as UV spectral slopes and optical to mid-infrared colors, and the possible differences between the two samples.

Finally through standard spectral fitting techniques we have determined the physical properties of the galaxies, such as total stellar mass, stellar ages and so on, and again we have studied the possible differences between the two samples.

Results. Our results indicate that LBG with $\text{Ly}\alpha$ in emission are on average a much younger and less massive population than the LBGs without $\text{Ly}\alpha$ emission. Both populations are forming stars very actively and are relatively dust free, although those with line emission seem to be even less dusty on average. We briefly discuss these results in the context of recent models for the evolution of Lyman break galaxies and $\text{Ly}\alpha$ emitters.

Key words. Galaxies:distances and redshift - Galaxies: evolution - Galaxies: high redshift - Galaxies: fundamental parameters -

1. Introduction

In the past years large samples of galaxies have been found up to the highest redshifts (Iye et al. 2006, Kashikawa et al. 2006, Bouwens et al. 2004) using techniques that rely on various color-selection criteria. Among the various methods, one of the most efficient is the Lyman break dropout technique (Steidel & Hamilton 1993) which is sensitive to the presence of the 912 Å break and is effective in finding star forming galaxies. This method requires a blue spectrum, implying low to moderate dust absorption. It was first designed to select ~ 3 galaxies (Steidel et al. 1996, Madau et al. 1996) with a typical magnitude limit $R > 25.5$, and then extended to higher redshift (Steidel et al. 1999; Ouchi et al. 2004; Giavalisco et al. 2004; Dickinson et al. 2004).

An alternative technique to find such distant objects is to search for $\text{Ly}\alpha$ emission, through very deep, narrow band imaging at selected redshift windows, as first shown by Cowie & Hu (1998). $\text{Ly}\alpha$ emitters (LAEs) are generally selected to have high restframe $\text{Ly}\alpha$ equivalent width, typically $EW \gtrsim 20\text{\AA}$, with no constraint on the continuum. Therefore, this method tends to select much fainter galaxies, compared to the general

LBGs population. Many $\text{Ly}\alpha$ emitters have now been found (e.g. Iye et al. 2006, Ouchi et al. 2004, Fujita et al. 2003), and several distant large scale structures or protoclusters have been discovered (e.g. Ouchi et al. 2005, Venemans et al. 2007).

Each of the two methods suffers from different selection bias: the two resulting galaxies population are partially overlapping, and it is not clear what is the relation between them.

Various scenarios have been proposed to explain the properties of $\text{Ly}\alpha$ emitters. Based on rest-frame optical photometry of LBGs, Shapley et al. (2001) concluded that LBGs with $\text{Ly}\alpha$ in emission are "old" (ages larger than $\sim 10^8$ yr), while "young" (ages less than ~ 100 Myrs) LBGs have $\text{Ly}\alpha$ in absorption. This could be explained if the young galaxies contain dust which absorbs the $\text{Ly}\alpha$ photons, while the older galaxies are more quiescent with less dust and superwinds which allow the $\text{Ly}\alpha$ photons to escape.

In alternative, other groups have suggested that strong $\text{Ly}\alpha$ emitters are instead young star forming galaxies, as derived from the blue colors and high equivalent widths of the $\text{Ly}\alpha$ emitters (e.g. Le Fevre et al. 1996, Malhotra & Rhoads 2002; Rhoads & Malhotra 2001; Tapken et al. 2004; Keel et al. 2002). Finally some authors have suggested that galaxies could have more than one $\text{Ly}\alpha$ bright emission phase (e.g. Thommes &

Meisenheimer 2005). An initial - primaevial - phase in which dust is virtually non-existent, and a later secondary phase in which strong galactic winds as observed in some Lyman break galaxies facilitate the escape of $\text{Ly}\alpha$ photons after dust has already been formed.

Clearly, it would be important to understand the real relation between galaxies with $\text{Ly}\alpha$ emission and the general LBG population, so that properties of the overall high redshift galaxy population, such as the total stellar mass density, can be better constrained.

To assess this issue we have analysed a sample of LBGs selected as B-dropouts from the GOODS-South sample, and with VLT spectroscopic confirmation (Vanzella et al. 2005, Vanzella et al. 2006). Given the good quality of the spectra it was possible to assess whether the galaxies have $\text{Ly}\alpha$ in emission or the line is absent and/or in absorption. We have then analysed how the observed properties and the derived physical properties such as total stellar mass and age depend on the nature of the $\text{Ly}\alpha$ emission.

All magnitudes are in the AB system (except where otherwise stated) and we adopt the Λ -CDM concordance cosmological model ($H_0 = 70$, $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$).

2. Sample and observational properties

From the GOODS-S public data survey we have selected all galaxies with spectroscopic redshift in the range $3.4 < z < 4.8$. These galaxies were initially selected as B-band (or V band for those at $z \geq 4.5$) dropouts by Giavalisco et al. (2004) (see this paper for the color selection criteria adopted) and with a z-band constrain $z < 26$. Spectroscopic observations were carried out with FORS2 in the frame of the GOODS project (Vanzella et al. 2005, Vanzella et al. 2006). The lower redshift cut is simply given by the actual availability of FORS2 spectra, given that the $\text{Ly}\alpha$ line falls at $\sim 5300 \text{ \AA}$, which is at the limit of efficiency for the observation-setup used by Vanzella et al. (2006) (FORS2 plus the 300I grism)¹; the highest redshift cut was chosen since at redshift higher than ~ 4.8 , the spectroscopic confirmations are almost exclusively based on the presence of the $\text{Ly}\alpha$ line in emission, while the objects with possible $\text{Ly}\alpha$ in absorption (or absent) become progressively more difficult to identify. In fact the most distant object identified with $\text{Ly}\alpha$ in absorption is at $z=4.788$.

From the public GOODS data we have retrieved the spectra, of quality A and B. The spectra with quality flag C were only included if our independently measured photometric redshift (from 14 bands photometry, Grazian et al. 2006) was in agreement with the spectroscopic redshift, within the errors. In the GOODS survey spectra are classified as emitters or absorbers, depending on the nature of the $\text{Ly}\alpha$ line. After removing AGNs, in the above redshift range there are 47 galaxies, of which 19 have the line in emission, and 28 have $\text{Ly}\alpha$ in absorption or absent. The implied ratio of line emitters/total is quite similar to the proportion of $\text{Ly}\alpha$ emitters that was found at redshift $z \sim 3$ in a large sample of LBGs by Steidel et al. (2000). We will call

the first sample LBG_L (LBG with Line emission) and the second sample LBG_N (LBGs with No line).

In Figure 1 (upper panel) we show the redshift distributions of the two samples, which are similar. From our GOODS-MUSIC database we have then gathered the complete 14 bands multicolor information, extending from the U band to the Spitzer $8\mu\text{m}$ band (for details of the catalog see Grazian et al. 2006). We have then studied the observational properties: in particular we have derived the restframe luminosity at $1400 \text{ \AA}$, the UV continuum slope β and the $z\text{--}[3.6]$ color which encompasses the $4000 \text{ \AA}$ break. The average observational properties of the two samples, as well as the physical properties that will be derived in the next section, are reported in Table 1, together with the uncertainties, derived using the standard deviation of the mean. In the last column of the Table we report, for each parameter, the probability value P given by the Kolmogorov-Smirnov test (KS-test) which tries to determine if the two datasets differ significantly. The KS-test has the advantage of making no assumption about the distribution of data.

The UV continuum slope was derived from the $i\text{--}z$ color, which spans a restframe wavelength range from $\sim 1500 \text{ \AA}$ to $\sim 1800 \text{ \AA}$ (at the mean redshift of the sample $z \sim 4$, and considering the central wavelengths of the ACS filters). Although the restframe wavelength baseline is not large (and perhaps a $i\text{--}J$ color would be preferred), we have used the $i\text{--}z$ color for two reasons: for a very few galaxies we do not have J band information available (see Grazian et al. 2006 for details); second, the UV slope derived in such way is more easily comparable to other UV slope values in the literature. Like other authors, we assume a standard power law spectrum with slope β ($f_\lambda \propto \lambda^\beta$, so that a spectrum that is flat in f_ν has $\beta = -2$). The $1216 \text{ \AA}$ break only starts to enter the i filter at $z > 4.7$ so we neglect the impact of neutral hydrogen absorption for all galaxies. The measured slopes are plotted in Figure 1 (lower panel) and range between -1 and -2.5 (with few exceptions). The uncertainties on the determined values range from 0.2 for the brightest objects to 0.8 for the fainter ones: in the figure we have indicated a median error on the individual values of 0.5 in the upper right corner. The average value is $\langle \beta \rangle = -1.8 \pm 0.13$ for the overall sample, and $\langle \beta \rangle = -2.0 \pm 0.11$ and -1.7 ± 0.13 considering the LBG_L and LBG_N separately. The UV slopes we find are consistent with the prediction of models for unobscured continuously star forming galaxies (e.g. Leitherer et al. 1999). They are also consistent with what found by other authors at a similar redshift: for example Overzier et al. (2006), for redshift $z=4.1$ LBGs and LAEs, find $\beta = -1.95$, Venemans et al. (2005) find $\beta = -1.65$ for $\text{Ly}\alpha$ emitters at $z=3.1$. The values are also similar to the average $\beta = -1.8 \pm 0.2$ of V606 dropouts in GOODS found by Bouwens et al. (2006).

On the other hand our values are a bit steeper than what found by Papovich et al. (2001) for LBGs at $z=3$ ($\langle \beta \rangle = -1.4$). Finally Shapley et al. (2003), found less steep values and claim a considerable difference between the LBG_L and LBG_N at $z \sim 3$, with the former having a steeper slope, with a positive dependence of slope on the $\text{Ly}\alpha$ equivalent width.

As can be seen in Figure 1, the values of the UV slopes for LBG_L and LBG_N are largely overlapping and a K-S does not reject the hypothesis that the two samples could be drawn from

¹ At lower redshift spectroscopic observations have been carried out with VIMOS but data are still not available to the public.

the same underlying distribution (the probability P is at $\sim 1\sigma$ see Table 1). However the median values are indeed somewhat different, and the trend is the same as what found by Shapley et al. (2003) given that the LBG_L are bluer on average than the LBG_N . The difference between our two samples ($\Delta\beta \sim 0.3$) is actually similar to that found by Shapley et al. (2003), who have a $\Delta\beta = 0.36$ between their most extreme groups i.e. the strong emitters and the strong absorbers (see Table 3 of the mentioned paper), but the significance of their $\Delta\beta$ is higher, given that they have a much larger sample with 800 objects in total, and therefore the uncertainties on the average β values are much lower than in our case. In conclusion, although we cannot claim such a strong dependence of β on the $\text{Ly}\alpha$ properties, we do indeed find a mild indication that LBG_L are bluer than LBG_N . To support this result, a similar difference is also found for LBG galaxies by Vanzella et al. (2006), who performed spectroscopy of B dropouts from the GOODS sample, and then stacking the spectra of galaxies with $\text{Ly}\alpha$ in emission and in absorption separately found that the former have a systematically bluer UV continuum than the latter.

We find no evidence for a slope-magnitude relation as found by other authors (e.g. Overzier et al. 2006) but the errors on the slope are quite large and the wavelength baseline for the slopes is rather small.

In Figure 2 we show the $z - [3.6]$ color plotted versus the total $[3.6]$ magnitude: this color encompasses the $4000\text{\AA}$ /Balmer breaks, and is sensitive both to dust and stellar ages. The $[3.6]$ magnitude corresponds to a rest-frame optical magnitude. The strong correlation between the optical flux and color is due to the magnitude cut at $z = 26$ that was used for the sample (see dotted line). It is clear from the figure that the LBG_N are on average both brighter (by ~ 0.7 magnitudes) at restframe optical wavelengths and redder (by more than half a magnitude) than the LBG_L . Given that all objects are relatively dust free (as inferred from the UV spectral slopes and consistently with the Lyman Break selection criteria), these differences indicate that the LBG_N have in general a more evolved stellar population and they are more massive. We will derive the physical properties in the next section.

3. Physical properties

3.1. Method

Using the multiwavelength GOODS-MUSIC data and the spectroscopic redshifts, we have then determined the physical properties of these galaxies, through a spectral fitting technique. The spectral fitting technique adopted here is the same that has been developed in previous papers (Fontana et al. 2003, Fontana et al. 2006), and similar to those adopted by other groups in the literature (e.g. Dickinson et al. 2003, Drory et al. 2004). Briefly, it is based on the comparison between the observed multicolor distribution of each object and a set of templates, computed with standard spectral synthesis models (Bruzual & Charlot 2003 in our case), and chosen to broadly encompass the variety of star-formation histories, metallicities and extinction of real galaxies. To compare with previous works, we have used the Salpeter IMF, ranging over a set of




Fig. 1. Upper panel: the redshift distribution of the two samples. Full black line are the LBG_L , light dashed line are the LBG_N . Lower panel: the UV slope calculated from the i-z colors, squares are the LBG_L and plus signs are the LBG_N .

metallicities (from $Z = 0.02Z_\odot$ to $Z = 2.5Z_\odot$) and dust extinction ($0 < E(B - V) < 1.1$, with a Calzetti or a Small Magellanic Cloud extinction curve). Details are given in Table 1 of Fontana et al. (2004). For each model of this grid, we have computed the expected magnitudes in our filter set, and found the best-fitting template with a standard χ^2 minimization. The stellar mass and other best-fit parameters of the galaxy, like SFR estimated from the UV luminosity and corrected for dust obscuration (with a typical correction factor of $A_V \sim 0.4$), age, τ (the star formation e-folding timescale), metallicity and dust extinction, are fitted simultaneously to the actual SED of the observed galaxy. The derivation of these parameters is explained in detail in the above paper and in Fontana et al. (2006), where also the uncertainties are discussed. In particular we note here that the stellar mass generally turns out to be the least sensitive to variations in input model assumptions, and the extension of the SEDs to the IRAC mid-IR data, tends to reduce considerably the formal uncertainties on the derived stellar masses. On the other hand, the physical parameter with highest associated uncertainty is the metallicity, given that the models are strongly degenerate when fitting broad band SEDs; a further limitation is that only 4 values were allowed in the modelling, i.e. 0.2, 0.02, 1 and 2.5 (respectively subsolar, solar and supersolar




Fig. 2. The $z-[3.6]$ color plotted vs. observed flux in $[3.6]$. Squares are the LBG_L and plus signs are the LBG_N .

metallicity), but in our samples none of the galaxies turned out to have a metallicity of 2.5.

3.2. Results

In Figure 3 we plot the distribution of the four basic physical parameters for the two samples, namely the total stellar mass, the star formation rate, the derived stellar age and the $E(B-V)$. In each figure we also indicate with an arrow the average values calculated for each sample.

Both the total stellar masses and the median ages are considerably lower for the LBG_L compared to the LBG_N . The average mass is $(5.0 \pm 1) \times 10^9 M_\odot$ for the LBG_L and $(2.3 \pm 0.8) \times 10^{10} M_\odot$ for the others, i.e. a factor of almost five higher. The K-S test gives a very low value, implying that the two populations are different from each other with $> 99.8\%$ probability. We can therefore conclude that the LBG_L are a factor of almost 5 less massive than the LBG_N .

The median ages are also quite different, with an average of 200 ± 50 Myr for the LBG_L , an age distribution that is very peaked towards small ages and is basically confined to values below 300 Myr. The LBG_N on the other hand have an average age of 410 ± 70 Myr (a factor ≥ 2 higher) and there are several galaxies with ages exceeding 1 Gyr, which is a considerable fraction of the cosmic time at redshift ~ 4 . Again performing a K-S test, the two populations are different with a probability $> 98\%$. We can conclude that the LBG_L are significantly younger galaxies compared to the LBG_N .

The star formation rates do not differ much for the two samples, with median values that are almost equal, $\langle SFR \rangle = 76 \pm 34 M_\odot \text{yr}^{-1}$ and $74 \pm 22 M_\odot \text{yr}^{-1}$ respectively and very similar distributions. The extinction values are slightly higher for the LBG_N with $\langle E(B-V) \rangle = 0.10 \pm 0.02$, while the LBG_L have $\langle E(B-V) \rangle = 0.07 \pm 0.01$. In both cases the K-S test does not reject the hypothesis that the 2 groups are drawn from the same population of galaxies, although the average $E(B-V)$ is higher for the LBG_N . Therefore both LBG_L and LBG_N are galaxies

with relatively little dust content, and significant star formation rates. In particular the average $E(B-V)$ is quite lower than the average values found by Shapley et al (2003) at redshift ~ 3 and Ouchi et al. (2004) at redshift ~ 4 but we find the same tendency for LBG_N to have somewhat higher dust content compared to LBG_L which is also supported by the difference in the UV spectral slopes discussed in the previous section.

The final two physical parameters that come as output from the modelling are the metallicity, and τ i.e. the star formation e-folding timescale. Each of them suffers from larger uncertainties as compared to e.g. the total stellar mass, as already discussed before. The mean values and the total distribution for these two parameters are very similar both for LBG_N and LBG_L , with the K-S test indicating that they are drawn from the same population. A more interesting and better constrained quantity to determine is the age/τ parameter that is an indication of the evolutionary state of the galaxies. It can be shown that passively evolving galaxies can be selected according to the physical criterion $age/\tau \geq 4$ (Grazian et al. 2006). All galaxies in our sample, both LBG_N and LBG_L have $age/\tau \leq 4$ as expected: however the difference in age discussed before is reflected also on the difference in age/τ , which is relatively larger for LBG_N with a mean value of $\langle age/\tau \rangle_{LBG_N} = 0.99$ while for LBG_L the mean value is 0.33. Furthermore all but one LBG_L have $age/\tau \leq 1$ while amongst LBG_N there are 7 objects with values between 1 and 4.

From this data we conclude that the LBG_L seem to be a population of younger and less massive galaxies. Given that all galaxies have relatively low dust content, there are some indication that LBG_L might be relatively less dusty than LBG_N in general, based on the results presented here and by other authors (Vanzella et al. 2006, Shapley et al. 2003 and Overzier et al. 2006). The star formation rates are high in all cases and are independent on the nature of the $\text{Ly}\alpha$ emission.

4. Discussion and Conclusions

The results obtained here seem to point towards a scenario where LBG_L are young and relatively small galaxies compared to the general LBGs population, are basically unobscured and are forming stars at very high rates. LBG_N on the other hand, are more massive, with masses as high as $10^{11} M_\odot$, and span a larger range of ages from few tens of Myrs to more than 1 Gyr. They are forming stars at comparable rates and are also relatively unobscured galaxies, although they appear slightly redder than LBG_L . This is therefore partially at variance with the only comparable large study, carried out at redshift ~ 3 by Shapley et al. (2003), who found a tendency for LBG_L to have in general higher ages, lower values of $E(B-V)$ and higher star formation rates. A very similar work by Iwata et al. (2005) on $z \sim 3$ LBGs from the Hubble Deep Field-South, also examined the relationship between spectroscopic features and parameters of the best-fit stellar population models in a sample of spectroscopically confirmed galaxies. They did not find any clear correlation between the $\text{Ly}\alpha$ emission, $\text{Ly}\alpha$ equivalent widths and the results of SED fitting. However the absence of any clear trend may be attributed to the very small number of sample galaxies in their study (13 in total).




Fig. 3. The physical properties of B and V-dropouts. Top left: distribution of total stellar mass; top right the extinction parameter $E(B-V)$ derived from the spectral fitting; middle left the total star formation rate derived from the spectral fitting; middle right: stellar ages; bottom left: τ the star formation e-folding time-scale; bottom right: the metallicity. Solid black and dashed light histograms denote LBG_L and LBG_N respectively. The black and light arrows indicate the mean values of each sample.

Some other observational studies seem to support our results: for example Overzier et al. (2006) find similar trends for a small sample of LBGs and LAEs in a $z=4.1$ protocluster, with indications that LAEs are young, dust free objects with blue UV optical colors and possibly less massive than UV selected LBGs with comparable star formation rates.

At higher redshift, Lai et al. (2006) constrained the ages of $\text{Ly}\alpha$ emitters at redshift 5 and found typical ages 50-100 Myr, i.e. comparable to our LBG_L . Finally at redshift 3, Gawiser et al. (2006) analysed a sample of narrow band selected $\text{Ly}\alpha$ emitters at $z \sim 3$ and concluded that they appear to have much less dust and lower stellar masses, compared to the generic LBG population at the same redshift. They were not able to constrain ages, which had a large range of allowed values. However given that they started from a narrow band sample, their objects are in general much fainter than the general LBGs population at the same redshift and clearly they are not selected to have the same broadband optical colors as the LBGs at the same redshift. In our case the optical colors and absolute restframe UV luminosity of the LBG_N and LBG_L are equal.

Our results give support to models that consider the $\text{Ly}\alpha$ emitters as precursors of spheroid galaxies, observed during the initial starburst phase, which is confined to a short period after its onset, due to the rapid formation of dust. Interestingly one such model (Thommes & Meisenheimer 2005) attempts to constrain the duration of this initial phase from observa-

tional data, finding $T_{\text{Ly}\alpha} \leq 350$ Myr. A very similar results was recently found by Mori & Umemura (2006) from simulation with a N body/hydrodynamics code: they follow the early stages of galaxy formation and find a first phase characterised by strong $\text{Ly}\alpha$ emission, which after 3×10^8 years quickly declines to fluxes below the observable level. The ages found in these models, are very similar to the range of allowed ages for our LBG_L as shown in Figure 3; on the other hand many of the LBG_N have ages exceeding 300 Myr. Clearly there are also very few LBG_L with larger ages and most of all, there are several LBG_N which are very young, with ages of a few tens of Myrs. The first few outliers could be easily accounted for given that the modelled physical parameter are obviously subject to uncertainties. However the very young LBG_N i.e. primeval objects without $\text{Ly}\alpha$ emission are hard to fit within the simple scenario proposed above. Probably more complex models in which the time scales for dust formation in primeval galaxies is variable and depends on other factors have to be invoked. In this context, several scenarios have been proposed, for example where the differential attenuation of the $\text{Ly}\alpha$ and continuum photons depends on the clumpiness of the medium (e.g. Hansen & Oh 2006), or where the timescales for the $\text{Ly}\alpha$ emission varies for galaxies residing in halos of different sizes, in the sense that in the most massive halos the $\text{Ly}\alpha$ luminosity declines abruptly after a much shorter time (10^7 ys) than in less

Table 1. Average properties of LBG_L and LBG_N

Property	LBG_L	LBG_N	P(K-S)
N_{gal}	19	28	
z	24.89 ± 0.13	24.85 ± 0.10	0.99
[3.6]	24.6 ± 0.2	23.8 ± 0.15	0.02
β	-2.0 ± 0.11	-1.7 ± 0.12	0.32
$L(1400)$ (cgs)	1.1×10^{29}	1.0×10^{29}	0.65
SFR ($M_\odot \text{yr}^{-1}$)	76 ± 34	74 ± 22	0.90
Age (Myrs)	200 ± 50	410 ± 70	0.02
E(B-V)	0.07 ± 0.01	0.10 ± 0.02	0.58
Mass ($M_\odot$)	$5 \pm 1 \times 10^9$	$2.3 \pm 0.8 \times 10^{10}$	<0.002
Metallicity ^a	0.55 ± 0.10	0.48 ± 0.08	1.0
τ	4.5 ± 1.4	5.5 ± 1.3	0.96
Age/ τ	0.33 ± 0.13	0.99 ± 0.23	0.06

Average observational and physical properties for the samples of LBG_L and LBG_N separately: β is the UV slope derived from the i-z color, $L(1400)$ is the total luminosity at 1400Åresframe; SFR is the total instantaneous star formation rate; Mass in the total stellar mass assembled in the galaxies; τ is the star formation e-folding timescale.

^a We remind that this parameter is the most uncertain from the SED fit output, and that only four values of metallicity were considered in the fitting procedure, see text for more details.

massive halos (Mao et al. 2007). However a full discussion of the models is beyond the scope of this paper.

Finally it also remains unclear whether the LBG_N are actually dustier than the LBG_L , which would be a natural consequence of this model: we do indeed find that LBG_L are slightly bluer than LBG_N (with the slope β determined directly from the observed colors) and consequently have on average lower E(B-V) parameter, as derived from the spectral fitting. However the differences are not very large. An important independent check would be to assess the dust content e.g. from far IR observations, rather than through the E(B-V) which is subject to larger uncertainties and is strongly degenerate with the fitted age.

References

Bouwens, R. J., et al. 2004, ApJ, 606, L25
 Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000
 Cowie, L. L., & Hu, E. M. 1998, AJ, 115, 1319
 Dickinson, M., Papovich, C., Ferguson, H.C., Budavari, T. 2003, ApJ, 587, 25
 Dickinson, M., et al. 2004, ApJ, 600, L99
 Drory, N., Bender, R., Feulner, G., et al. 2004, ApJ, 608, 742
 Fontana, A., Salimbeni, S., Grazian, A., et al. 2006 A&A in press
 Fontana, A., et al. 2004, A&A, 424, 23
 Fontana, A., et al. 2003, ApJ, 594, L9
 Fujita, S. S., et al. 2003, AJ, 125, 13
 Gawiser, E., et al. 2006, ApJ, 642, L13
 Giavalisco, M., et al. 2004, ApJ, 600, L103
 Grazian, A., et al. . 2007 A&A accepted astroph/0701233
 Grazian, A., Fontana, A., De Santis, C., Nonino, M., Salimbeni, S., Giallongo, E., Cristiani, S., Gallozzi, S. and Vanzella, E. 2006a A&A 449, 951.
 Hansen, M., & Oh, S. P. 2006, MNRAS, 367, 979
 Iye, M., et al. 2006, Nature, 443, 186
 Iwata, I., Inoue, A. K., & Burgarella, D. 2005, A&A, 440, 881
 Kashikawa, N., et al. 2006, ApJ, 648, 7

Keel, W. C., Wu, W., Waddington, I., Windhorst, R. A., & Pascarelle, S. M. 2002, AJ, 123, 3041
 Lai, K. et al. 2006, Accepted for publication in ApJ astro-ph/0610572
 Le Fevre, O., Deltorn, J. M., Crampton, D., & Dickinson, M. 1996, ApJ, 471, L11
 Leitherer, C., et al. 1999, ApJS, 123, 3
 Madau, P., Ferguson, H. C., Dickinson, M. E., Giavalisco, M., Steidel, C. C., & Fruchter, A. 1996, MNRAS, 283, 1388
 Malhotra, S., & Rhoads, J. E. 2002, ApJ, 565, L71
 Mao, J., Lapi, A., Granato, G. L., de Zotti, G., & Danese, L. 2006, ArXiv Astrophysics e-prints, arXiv:astro-ph/0611799
 Mori, M., & Umemura, M. 2006, Nature, 440, 644
 Ouchi, M., et al. 2005, ApJ, 620, L1
 Ouchi, M., et al. 2004, ApJ, 611, 660
 Overzier, R. et al. 2006, ApJ submitted, astro-ph/0601223
 Papovich, C., Dickinson, M., & Ferguson, H. C. 2001, ApJ, 559, 620
 Rhoads, J. E., & Malhotra, S. 2001, ApJ, 563, L5
 Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, ApJ, 588, 65
 Shapley, A. E., Steidel, C. C., Adelberger, K. L., Dickinson, M., Giavalisco, M., & Pettini, M. 2001, ApJ, 562, 95
 Steidel, C. C., & Hamilton, D. 1993, AJ, 105, 2017
 Steidel, C. C., Giavalisco, M., Pettini, M., Dickinson, M., & Adelberger, K. L. 1996, ApJ, 462, L17
 Steidel, C. C., Adelberger, K. L., Shapley, A. E., Pettini, M., Dickinson, M., & Giavalisco, M. 2000, ApJ, 532, 170
 Tapken, C., Appenzeller, I., Mehlert, D., Noll, S., & Richling, S. 2004, A&A, 416, L1
 Thommes, E., & Meisenheimer, K. 2005, A&A, 430, 877
 Vanzella, E., et al. 2006, Proceedings for the conference "At the Edge of the Universe: Latest results from the deepest astronomical surveys", astroph/0612182
 Vanzella, E., et al. 2006, A&A, 454, 423
 Vanzella, E., et al. 2005, A&A, 434, 53
 Venemans, B. P., et al. 2007, A&A, 461, 823
 Venemans, B. P., et al. 2005, A&A, 431, 793